

Nutritional requirements

The composition of medium for the tissue culture is the most important key factor in the successful culture of plant cells. The medium should be accurately defined of inorganic and organic chemical additives so as to provide i) the nutrients for the survival of the plant cells, tissues and organs under culture and ii) the optimal physical condition of pH, osmotic pressure, *etc.*

In the culture of plant cells formulating optimum type of medium favorable for *in vitro* culture was achieved many years ago. The **Knop's** (1865) mineral solution was the widely used medium by early investigators. **Gautherat** (1939) developed callus culture medium from **Uspenski** and **Uspenskaia** (1925) nutrient solution. A systematic study of mineral requirements of plant tissue and organs in culture was made by **Murashige** and **Skoog** (1962) followed by the scientists **Linsmaier** and **Skoog** (1965), **Vasil** and **Hildebrandt** (1966) and **Nitsch** and **Nitsch** (1969) resulting in several media to suit particular needs.

Nutrients

A standard basal medium consists of a balanced mixture of macronutrients and micronutrients (usually salts of chlorides, nitrates, sulphates, phosphates and iodides of Ca, Mg, K, Na, Fe, Zn and B, a carbon source, vitamins, phytohormones and organic additives. Among the above mentioned nutrients some are essential and some are optional. The essential components include inorganic nutrients and organic nutrients like carbohydrates besides phytohormones and vitamins, organic additives like natural extract and liquid endosperm are optional.

Inorganic salts

Inorganic nutrients of a plant cell culture are those required by the normal plants. The optimum concentration of each nutrient for achieving maximum growth rates varies considerably. The major elements are N, P, K, S, Mg and Ca. Other nutrients such as Co, Fe, B, Zn, Mo, Cu, I are microelements.

Macroelements

Nitrogen

Of all the mineral nutrients N plays a vital role in growth and differentiation of cultured tissues. The range of inorganic nitrogen varies from 25 mM to 60 mM according to the requirements. Nitrogen is generally supplied in the form of NH_4 along with NO_3 .

Ammonium ion as nitrogen source is usually unsuitable, probably because under such conditions the pH of the medium has a tendency to fall below 5 during culture, resulting in reduced availability of nitrogen. Cells can be grown with NH_4 as the sole N source when the medium is provided with organic acids such as malate, succinate, citrate or fumarate. Further, the concentration of $\text{NH}_4\text{-N}$ should not exceed 8 mM. Generally $\text{NO}_3\text{-N}$ can be used as a sole N source but often there is a beneficial effect if the media contains $\text{NH}_4\text{-N}$.

Phosphorus

Phosphorus is usually supplied in the form of phosphates. It is the primary buffering constituent in tissue culture media. Phosphorus levels greater than 2mM are often inhibitory to growth of tissues.

Potassium

The optimum concentration of K needed is 20 mM. At low nitrogen concentration presence of potassium enhances the formation of somatic embryos. The medium supplemented with potassium nitrate produces more embryos than the medium with ammonium nitrate.

Sulphur

Sulphur is provided in the form of sulphates. Besides, the sulphur containing amino acids like L-cysteine, L-methionine and glutathione are satisfactory sources for sulphur.

Calcium and Magnesium

The optimal concentration of Ca required is 3mM. An antagonism between Ca and Mg has been demonstrated and it was found that an increase in the concentration of one element increased the requirement for the other.

Microelements

The microelements *viz.*, Fe, Mn, B, Zn, Mo, Cu, I and Co have a profound effect on growth of tissue *in vitro*. The availability of the iron is reduced at high pH due to precipitation. To avoid this, Fe is supplied as chelated EDTA complex. These elements produce toxic effect, if they are applied at higher level. A good growth of tissue can be achieved when the concentration of microelements was reduced to 10 per cent of the original level.

Organic nutrients

Carbohydrates

Carbohydrates are used as carbon sources. The standard carbon source is sucrose at a concentration of 2-5 per cent. Monosaccharides like glucose or fructose can also be used as carbon sources but are generally less suitable. Sucrose is the best source, since sucrose is dehydrolysed into usable sugars during autoclaving.

Vitamins

Vitamins are supplemented with medium to achieve the best growth of the tissues. Among the vitamins only thiamine HCL (B₁) seems to be universally required. Other vitamins are pyridoxine HCL (B₆), nicotinic acid (B₃) and calcium pantothenate (B₅). Specific requirement of each one varies with the plant species subject to culture.

Phytohormones

These are organic compounds, other than nutrients, which influence growth, differentiation and multiplication. They required in very minute quantity in the media. The requirement for these substances varies considerably with the tissue and it also depends on their endogenous level. There are many commercially available synthetic substances that mimics the PGR specific to certain species. Testing of various types, concentrations and mixtures of growth substances during the development of a tissue culture protocol for a new species is essential before using a new PGR in plant tissue culture.

There are different groups of PGRs commonly used in the media. They are auxins, cytokinins, gibberellins, ethylene and abscisic acid. Additional substances gaining recognition as hormones in plant tissue culture are: polyamines, jasmonates, salicylic acid and brassinosteroids.

1. Auxin

In nature, the hormones of this group are involved with elongation of stem, internodes, tropism, apical dominance, abscission, rooting etc. In tissue culture auxins have been used for cell division and root differentiation. The commonly used auxins in tissue culture are

1. Indole-3-acetic acid (IAA)
2. Indole-3-butyric acid (IBA)
3. Naphthalene acetic acid (NAA)
4. Dichlorophenoxyacetic acid (2, 4-D)

Auxins are usually dissolved in either ethanol or dilute NaOH.

2. Cytokinins

These hormones are essential for cell division, modification of apical dominance, shoot differentiation etc. In tissue culture media, cytokinins are incorporated mainly for cell division, differentiation of adventitious shoots from callus and organ & shoot proliferation.

Commonly used cytokinins are


1. Benzylamino purine (BAP)
2. Isopentenyl adenine (2-ip)
3. Furfurylamino purine (kinetin)
4. Zeatin

Cytokinins are generally dissolved in dilute HCl or NaOH.

Auxin - Cytokinin Interaction

1. High auxin and low cytokinin ratio: Initialize root formation, embryogenesis and callus formation.
2. Low auxin and high cytokinin ratio: Induce formation of adventitive or axillary shoots.
3. The auxin-cytokinin ratio is also essential for chloroplast formation and other processes.

Effect of different auxin and cytokinin concentration on tissue development



3. Gibberellins

Naturally occurring plant hormones involved in internode elongation, enhancement of flower, fruit and leaf size, germination and vernalization in plants. Among the 20 known gibberellins, GA₃ is used widely. Compared to auxins and cytokinins, gibberellins are used very rarely. They stimulate normal development of plantlets from *in vitro* formed adventitious embryos. They are soluble in cold water.

4. Ethylene

A gaseous plant hormone involved in fruit maturation, abscission, and senescence. All kinds of plant tissue cultures produce ethylene and the rate of production increases under stress conditions. Use of ethylene precursor (2-chloroethylphosphonic acid) in tissue culture may be promotory or inhibitory for the same process in different species. For example, it promoted somatic embryogenesis in *Zea mays* whereas the same process was inhibited in *Hevea brasiliensis*.

5. Abscisic acid

A plant hormone involved in abscission, enforcing dormancy and regulating early stages of embryo development. It is required for normal growth and development of somatic embryos and promotes morphogenesis.

6. Brassinosteroids

It promotes shoot elongation at low concentrations and strongly inhibits root growth and development. It also promotes ethylene biosynthesis and epinasty.

7. Jasmonates

Jasmonates are represented by jasmonic acid and it is a methyl ester. Jasmonic acid is considered to be a new class of plant growth substance. It inhibits many processes such as embryogenesis, seed germination, pollen germination, flower bud formation, chlorophyll formation. It is involved in differentiation, adventitious root formation, breaking of seed dormancy and pollen germination.

8. Polyamines

There is some controversy as to whether these compounds should be classified with hormones. They appear to be essential in growth and cell division.

9. Salicylic acid

It is thought to be a new class of plant growth substances. It promotes flowering, inhibits ethylene biosynthesis and reverses the effects of ABA.

Organic additives

Amino acids like glutamine, asparagine and nitrogen base like adenine are used as additives in tissue culture media. The organic acids citrate, malate, succinate and fumarate are used when the medium has nitrogen in ammoniacal form.

A wide variety of complex natural extracts like coconut water (liquid endosperm) tomato and orange juices is also used during media preparation. These complex substances possess a number of amino acids, vitamins, sugars, sugar alcohols, growth regulators and other unidentified substances with growth promoting qualities. However, these should be avoided because of their unknown and variable composition. Among the natural extract coconut water is widely used as a source of cytokinin and various amino acids.

Other complex substances often used in tissue culture media are: casein hydrolysate, yeast extract and malt extract. Potato extract is also used in China for cereal anther cultures.

Physical form of media

I. Solidified medium

Advantages

1. Explants are easily seen and recovered.
2. No Special aeration required.
3. Shoots grow more orderly.
4. Long term maintenance is possible.
5. Use of simple containers occupy little space

Disadvantages

1. Slower rate of multiplication.
2. Limited surface of explant is in contact with the medium.

II. Liquid medium

Advantages

1. Callus easily break up and shed as cells to establish a fine suspension.

2. Faster rate of multiplication
3. A greater surface of explants is in contact with medium.
4. Toxic metabolites will effectively be dispersed.

Disadvantages

1. Recovery is difficult.
2. Growth is disoriented.
3. Seeds would not germinate.
4. Protocorms and plantlets become brown and dry. Seeds submerged will show vitrification.

Commonly used tissue culture media

1. **MS** (Murashige and Skoog, 1962) and **LS** (Linsmaier and Skoog, 1972) media are used for regeneration of both monocots and dicots.
2. **B5** (Gamborg *et al.*, 1969) developed for culture of soybean cell suspensions but also has been effectively used for variety of plant regeneration. **B5** and its various derivatives have been valuable for cell and protoplast cultures.
3. **SH**: Schenk and Hildebrandt (1972) introduced this for culture of monocots and dicots. Widely used for legumes.
4. **WPM**: Lloyd and McCown (1980-1981). This is post MS media. WPM is increasingly used for propagation of Ornamental shrubs and trees in commercial labs.
5. **N6** was developed by Chu for cereal anther culture and also used for anther culture of other species.
6. In special cases, NN (Nitsch and Nitsch) was also used.

The physical form of a tissue culture medium, like the combination of nutrients is more important, since the uptake of nutrients by the tissues, their growth and development are dependent on it. To maintain the physical form of medium suitable for culturing, care should be taken to maintain the necessary (1) hydrogen ion concentration. (2) gelling agent and (3) osmotic pressure of the medium.

Hydrogen ion concentration (pH)

The pH of the medium is usually adjusted between 5.0 and 6.0 prior to the addition of agar and autoclaving. The extremes of pH should be avoided as this will block the availability of some of nutrients to the inoculum. A pH of 5.8 is found to be optimum for plant tissue

culture. Generally, pH higher than 6.0 give a very hard medium and a pH below 5.0 do not allow satisfactory solidification of agar. Further, pH in media changes during growth of plant tissues and this drift in pH is comparatively low in media with high salt concentration, because of their greater buffering capacity.

Gelling agents

Generally tissue culture media are solidified with any of the gelling agents. Agar is widely used for solidification of the medium. The optimum concentration of agar used ranges from 0.8-1.0 per cent (W/V). If the concentration of the agar is increased, the medium becomes hard and does not allow the diffusion of nutrient into the tissues. Gelatin, silica gel, acryl amide gel and starch copolymers are also used as substitutes for agar. Sometimes, the solid media will accumulate the toxic substances namely, oxidised, phenolic compounds released from tissue and hamper further growth of tissue. To absorb the toxic substances, 1 percent activated charcoal is added to the medium. One disadvantage of adding activated charcoal is that it would adsorb the growth regulators.

Liquid medium (the medium without any gelling agent) is suitable for suspension culture and it is superior to other media for the following reasons: 1) does not have impurities as in the agarified medium where the agar contains impurities, 2) aeration can be provided to the cells by keeping the suspensions under constant shakings and 3) toxic substances released from the tissues will not accumulate or localize; the substances get diluted. In liquid media cultures, filter paper bridges or glass wool can be used to support culture tissue.

Osmotic pressure

The cell cultured *in vitro* is mostly osmotically fragile and hence the osmotic pressure of the medium should be maintained at optimal level. This problem is serious one when liquid media are used. To adjust the osmotic pressure, stabilizers otherwise known as osmoticums *viz.*, sorbitol and mannitol (sugar alcohol) are used. These are non metabolisable sugars. The soluble sugar like sucrose, fructose, galactose, *etc.*, is also effective. The sucrose is added to the medium not only to provide energy but also to maintain a suitable osmolarity in the medium.

Several modifications have been made in the basic media evolved for various types of plant tissue cultures and the modifications are ever continuing processes in the field of plant

tissue culture. The reason behind this is that the selection of a particular culture medium for a particular species is difficult. Considering the difficulties the following approaches may be taken into consideration to identify a suitable medium for the work. 1) literature survey for work on similar objectives or near relative species and to try out the media in the reports, 2) experimentation with several of the well known media incorporating some variables and 3) conducting broad spectrum experiments involving most of the components (minerals, carbon source and phytohormones) with different treatments. The suitable combinations can be identified when the desired response is achieved.

Environmental factors influencing plant tissue culture

1. Genotype or variety of the plant material
2. Explant selection and its size.
3. Medium:
 - (i) Nutrients,
 - (ii) Growth regulators and
 - (iii) Other additives
4. Culture Environment:
 - (i) Temperature,
 - (ii) Relative Humidity (RH) and
 - (iii) Light

Questions

1. The mineral solution widely used by early investigators was

- a) **Knop's medium**
- b) MS medium
- c) White's medium
- d) None of the above

2. The medium in tissue culture should provide

- a) Nutrients for the survival of the plant cells, tissues and organs under culture
- b) Optimal physical condition of pH, osmotic pressure, *etc*
- c) **Both a & b**
- d) None of the above

3. The callus culture was first developed by

- a) Skoog
- b) White
- c) **Gautherat**
- d) None of the above

4. The systematic study of mineral requirements of plant tissue and organs in culture was made by

- a) Murashige and Skoog
- b) Vasil and Hildebrandt
- c) Nitsch and Nitsch
- d) **All the above**

5. The mineral nutrient that plays a vital role in growth and differentiation of cultured tissues is

- a) **N**
- b) P
- c) K
- d) All the above

6. Nitrogen is generally supplied in form.

- a) **NH₄**
- b) NO₃
- c) NO₂
- d) All the above

7. The range of inorganic nitrogen varies between in the tissue culture medium.

- a) **25 mM - 60 mM**
- b) 250 mM - 600 mM
- c) 2.5 mM – 6.0 mM
- d) 0.25 mM – 0.60 mM

8. Phosphorus levels greater than are inhibitory to growth of tissues.

- a) 2 mM
- b) 20 mM
- c) 200 mM
- d) 0.2 mM

9. The optimum concentration of K needed is

- a) 2 mM
- b) 20 mM**
- c) 200 mM
- d) 0.2 mM

10. At low nitrogen concentration presence of element enhances the formation of somatic embryos

- a) K**
- b) P
- c) Ca
- d) Mg

11. Sulphur is provided in the form of sulphates

- a) Sulphates**
- b) Sulphites
- c) Both a & b
- d) None of the above

12. The standard carbon source is

- a) Sucrose (2-5 per cent)**
- b) Glucose
- c) Fructose
- d) None of the above

13. The standard carbon source is

- a) Sucrose (2-5 per cent)**
- b) Glucose
- c) Fructose
- d) None of the above

14. Universally required vitamin in tissue culture medium is

- a) Thiamine HCL**
- b) Pyridoxine HCL
- c) Nicotinic acid
- d) Calcium pantothenate

15. Vitamin (s) used in tissue culture medium is/are

- a) Calcium pantothenate
- b) Pyridoxine HCL
- c) Nicotinic acid
- d) All the above**

16. The commonly used auxins in tissue culture are

- a) IAA, IBA, NAA, 2,4 D**
- b) BAP, 2-ip, Kinetin, Zeatin

c) GA₃

d) All the above

17. The commonly used cytokinins in tissue culture are

a) IAA, IBA, NAA, 2,4 D

b) BAP, 2-ip, Kinetin, Zeatin

c) GA₃

d) All the above

18. The auxins are diluted in

a) Ethanol

b) Dilute NaOH

c) Both a & b

d) None of the above

19. The cytokinins are diluted in

a) Dilute HCl

b) Dilute NaOH

c) Both a & b

d) None of the above

20. In tissue culture auxins have been used for

a) Cell division and root differentiation

b) Differentiation of adventitious shoots from callus and organ & shoot proliferation

c) Normal development of plantlets from *in vitro* formed adventitious embryos

d) None of the above

21. In tissue culture cytokinins have been used for

a) Cell division and root differentiation

b) Differentiation of adventitious shoots from callus and organ & shoot proliferation

c) Normal development of plantlets from *in vitro* formed adventitious embryos

d) None of the above

22. In tissue culture gibberellins have been used for

a) Cell division and root differentiation

b) Differentiation of adventitious shoots from callus and organ & shoot proliferation

c) Normal development of plantlets from *in vitro* formed adventitious embryos

d) None of the above

23. High auxin and low cytokinin ratio favours

- a) **Initialize root formation, embryogenesis and callus formation** b) Induce formation of adventitive or axillary shoots
c) Chloroplast formation and other processes d) None of the above

24. Low auxin and high cytokinin ratio favours

- a) Initialize root formation, embryogenesis and callus formation **b) Induce formation of adventitive or axillary shoots**
c) Chloroplast formation and other processes d) None of the above

25. The auxin-cytokinin ratio is essential for

- a) Initialize root formation, embryogenesis and callus formation b) Induce formation of adventitive or axillary shoots
c) Chloroplast formation and other processes **d) All the above**

26. The naturally occurring gibberellin is involved in

- a) Internode elongation b) Enhancement of flower, fruit and leaf size
c) Germination and vernalization **d) All the above**

27. The gibberellins are soluble in

- a) Dilute HCl b) Dilute NaOH
c) Cold water d) Hot water

28. Ethylene is involved in

- a) Fruit maturation b) Abscission
c) Senescence **d) All the above**

29. Abscissic acid is involved in

- a) Enforcing dormancy b) Abscission
c) Regulating early stages of embryo development **d) All the above**

30. Abscissic acid is required for

- a) Normal growth b) **Promotes morphogenesis**
c) Development of somatic embryos **d) All the above**

31. promotes ethylene biosynthesis and epinasty.

- a) **Brassinosteroids**
- b) Jasmonates
- c) Polyamines
- d) Salicyclic acid

32. Jasmonic acid is a

- a) **Methyl ester**
- b) Ethyl ester
- c) Ether
- d) None of the above

33. The optimum pH of the tissue culture is

- a) **5.8**
- b) 6.0
- c) 4.0
- d) 7.0

34. The substitutes for agar are

- a) Gelatin
- b) Silica gel
- c) Acryl amide gel
- d) **All the above**

35. The osmoticums used in tissue culture medium are

- a) Sorbitol and mannitol
- b) Sucrose
- c) Fructose and galactose
- d) **All the above**