


## Soil aggregates-significance-soil consistency-soil crusting

### Soil aggregates

Soil aggregates are 'clumps' of soil particles that are held together by moist clay, organic matter (such as roots), by organic compounds (from bacteria and fungi) and by fungal hyphae (*pronounced "highfee"*). Aggregates vary in size from about 2 thousandths of a millimetre across up to about 2 millimetres across, and are made up of particles of varying sizes. Some of these particles fit closely together and some do not and this creates spaces of many different sizes in the soil. These spaces, or pores, within and between soil aggregates are essential for storing air and water, microbes, nutrients and organic matter. Soils with many aggregates are called "well-aggregated". Such soils are more stable and less susceptible to erosion.

There are two ways that bacteria could be involved in soil aggregation. One way is by producing organic compounds called polysaccharides. Bacterial polysaccharides are more stable than plant polysaccharides, resisting decomposition long enough to be involved in holding soil particles together in aggregates. The other way bacteria are involved in soil aggregation is by developing a small electrostatic charge that attracts the electrostatic charge on clay surfaces, bringing together small aggregates of soil.



Fungi grow in long, threadlike structures, called hyphae. The amount of aggregation in the soil has been found to relate to the length of fungal hyphae in the soil. Fungi help to form aggregates in the soil by enmeshing soil particles with their hyphae and forming cross-links between soil particles. Mycorrhizal fungi and fungi that colonise fresh organic matter

are believed to be the most important for assisting with stabilisation of soil particles into aggregates.

### **Significance of soil aggregation**

Crop growth is often constrained by poor root development, by slow water infiltration and water movement through the soil, and by poor soil aeration. These constraints are often associated with poor soil porosity. Soil aggregation is important to developing and maintaining good soil porosity and hence to good root growth and to movement of soil water and gases. With more soil in water stable aggregates, it is expected that:

- The rate of water infiltration and percolation will increase
- Soil crusting will be less – which improved root penetration and access to soil moisture and nutrients and emergence of seedlings
- Resistance to the splash effect of raindrops will increase and soil erodibility will decrease; and
- Runoff will decrease, making more water available to the crop.

The importance of soil physical properties to crop growth, including soil aggregation properties, is often under-estimated by producers, and practical exercises can be useful in conveying information on the importance of these properties.

### **Soil Consistence**

**Soil consistence** is defined as “the resistance of a soil at various moisture contents to mechanical stresses or manipulations”.

It combines both the ‘**cohesive**’ and ‘**adhesive**’ forces, which determine the ease with which a soil can be reshaped or ruptures.

#### **Adhesion**

Molecular attraction that holds the surfaces of two substances (eg. Water and soil particles) in contact

#### **Cohesion**

Holding together: force holding a solid or liquid together, owing to attraction between like molecules. Decreases with rise in temperature

Soil consistence is described at three moisture levels namely ‘wet’, ‘moist’ and ‘dry’.

**1. Wet soils:** Consistency is denoted by terms stickiness and plasticity

**Stickiness** is grouped into four categories namely i) non sticky, ii) slightly sticky, iii) sticky and iv) very sticky

**Plasticity** of a soil is its capacity to be moulded (to change its shape depending on stress) and to retain the shape even when the stress is removed. Soils containing more than about 15% clay exhibit plasticity – pliability and the capacity of being molded. There are four degrees in plasticity namely i) non plastic, ii) slightly plastic, iii) plastic and iv) very plastic.

**2. Moist soil:** Moist soil with least coherence adheres very strongly and resists crushing between the thumb and forefinger. The different categories are

- i. Loose-non coherent
- ii. Very friable - coherent, but very easily crushed
- iii. Friable - easily crushed
- iv. Firm - crushable with moderate pressure
- v. Very firm - crushable only under strong pressure
- vi. Extremely firm - completely resistant to crushing. (type and amount of clay and humus influence this consistency)

**3. Dry soil:** In the absence of moisture, the degree of resistance is related to the attraction of particles for each other. The different categories are

- i) Loose - non coherent
- ii) Soft - breaks with slight pressure and becomes powder
- iii) Slightly hard - break under moderate pressure
- iv) Hard - breaks with difficulty with pressure
- v) Very hard - very resistant to pressure
- vi) Extremely hard - extreme resistance and cannot be broken

## Soil crusting

## **Formation mechanism:**

Soil crusts usually are formed as a result of compaction at the immediate surface due to an externally applied force. This force is supplied primarily by the impact of raindrops as the soil is wetted and the radiant energy of the sun as the soil dries. When the rain drops fall on dry soil.

## **Bulk density and particle density of soils & porosity**

### **Particle Density**

The weight per unit volume of the solid portion of soil is called particle density. Generally particle density of normal soils is 2.65 grams per cubic centimeter. The particle density is higher if large amount of heavy minerals such as magnetite, limonite and hematite are present in the soil. With increase in organic matter of the soil the particle density decreases. Particle density is also termed as true density.

Table Particle density of different soil textural classes

Textural class	Particle density ( $\text{g}/\text{cm}^3$ )
Coarse sand	2.655
Fine sand	2.659
Silt	2.798
Clay	2.837

### **Bulk Density**

The oven dry weight of a unit volume of soil inclusive of pore spaces is called bulk density. The bulk density of a soil is always smaller than its particle density. The bulk density of sandy soil is about  $1.6 \text{ g}/\text{cm}^3$ , whereas that of organic matter is about 0.5. Bulk density normally decreases, as mineral soils become finer in texture. The bulk density varies indirectly with the total pore space present in the soil and gives a good estimate of the porosity of the soil. Bulk density is of greater importance than particle density in understanding the physical behavior of the soil. Generally soils with low bulk densities have favorable physical conditions.

### **Bulk density of different textural classes**

<b>Textural class</b>	<b>Bulk density (g/cc)</b>	<b>Pore space (%)</b>
Sandy soil	1.6	40
Loam	1.4	47
Silt loam	1.3	50
Clay	1.1	58

## Factors affecting bulk density

### 1. Pore space

Since bulk density relates to the combined volume of the solids and pore spaces, soils with high proportion of pore space to solids have lower bulk densities than those that are more compact and have less pore space. Consequently, any factor that influences soil pore space will affect bulk density.

### 2. Texture

Fine textured surface soils such as silt loams, clays and clay loams generally have lower bulk densities than sandy soils. This is because the fine textured soils tend to organize in porous grains especially because of adequate organic matter content. This results in high pore space and low bulk density. However, in sandy soils, organic matter content is generally low, the solid particles lie close together and the bulk density is commonly higher than in fine textured soils.

### 3. Organic matter content

More the organic matter content in soil results in high pore space there by shows lower bulk density of soil and vice-versa.

## **Porosity**

Soil porosity refers to that part of a soil volume that is not occupied by soil particles or organic matter.

The pore space of a soil is the space occupied by air and water. The amount or ratio of pore space in a soil is determined by the arrangement of soil particles like sand, silt and clay. In sandy soils, the particles are arranged closely and the pore space is low. In clay soils, the particles are arranged in popous aggregates and the pore space is high. Presence of organic matter increases the pore space.

### **Factors influencing pore space**

#### **Soil texture**

Sandy surface soil : 35 to 50 %

Medium to fine textured soils : 50 to 60 %

Compact sub soils : 25 to 30%

#### **Crops / vegetation**

Some crops like blue grass increases the porosity to 57.2% from the original 50%

Cropping reduces the porosity as cultivation reduces the organic matter content and hence decrease in granulation. Virgin soils have more pore space.

Continuous cropping reduces pore space than intermittent cropping. More the number of crops per year, lesser will be the pore space particularly macro pores.

Conservation tillage and no tillage reduces porosity than conventional tillage

#### **Size of pores**


1. Macro pores (non-capillary pores) : diameter >0.05 mm

2. Micro pores (capillary pores) : diameter < 0.05 mm

In macro pores, air and water moves freely due to gravitation and mass flow. In micro pores, the movement of air and water is very slow and restricted to capillary movement and diffusion.

Sandy soil have more macro pores and clay soils have more micro pores. So in sandy soils, water and air movement is rapid due to macro pores though the pore space is higher and in clay soils the air and water is slower due to micro pores though the total pore space is higher.

Loamy soils will have 50% porosity and have equal portion of macro and micro pores.



### Significance and manipulation of soil porosity

The bulk density and pore space are inter related. Development of low bulk density values also means the development of large amount of pore spaces. In nature, low bulk density values are usually found in soils with high organic matter contents. High biological activities are necessary for formation and large accumulation of organic matter. Together with the effect of soil organisms, the high humus content will encourage aggregation, increasing in this way soil porosity, and thereby decreasing bulk density values. The cultivation effect of the

soil macro and micro fauna produces an intricate system of macropores, which is a major factor for lowering the bulk density of soil. Continuous cropping is noted to decrease the amount of organic matter in soils, and is expected to decrease soil aggregation. Tillage by ploughing is designed to increase the pore space in soils, but is in fact decreasing organic matter. To alleviate these problems conservation tillage and no tillage have been introduced. Though many claimed that this increased the organic matter, the latter have not always increased the total pore space.

\*\*\*\*\*